

15-819 Homotopy Type Theory

Week 2 Lecture Notes

Clive Newstead and Enoch Cheung

9/16/2013 and 9/18/2013

Foreword

These will undergo substantial revision and expansion in the coming week.

Recall from last time that we can think of the judgement A **true** as meaning ‘ A has a proof’ and of A **false** as ‘ A has a refutation’, or equivalently ‘ $\neg A$ has a proof’. These atomic judgements give rise to hypothetical judgements of the form

$$A_1 \text{ true}, A_2 \text{ true}, \dots, A_n \text{ true} \vdash A \text{ true}$$

The inference rules of **intuitionistic propositional logic** then give rise to the structure of a Heyting algebra, called the *Lindenbaum algebra*.

1 Lindenbaum algebras

Recall that **IPL** has the structure of a preorder, where we declare $A \leq B$ if and only if $A \text{ true} \vdash B \text{ true}$. Let T be some theory in **intuitionistic propositional logic** and define a relation \simeq on the propositions in T by

$$A \simeq B \quad \text{if and only if} \quad A \leq B \text{ and } B \leq A$$

The fact that \simeq is an equivalence relation follows from the more general fact if (P, \leq) is a preorder and a relation \equiv is defined on P by declaring $p \equiv q$ if and only if $p \leq q$ and $q \leq p$, then \equiv is an equivalence relation on P .

Definition. The *Lindenbaum algebra* of T is defined to be the collection of \simeq -equivalence classes of propositions in T . Write $A^* = [A]_{\simeq}$. The ordering on the Lindenbaum algebra is inherited from \leq .

Theorem. The judgement $\Gamma \vdash A \text{ true}$ holds if and only if $\Gamma^* \vdash A^*$ holds in every Heyting algebra.

Proof. Exercise. □

2 Decidability and stability

Definition. A prop is *decidable* if and only if $A \vee \neg A \text{ true}$.

Decidability is what separates constructive logic from classical logic: in classical logic, every proposition is decidable (this is precisely the **law of the excluded middle**), but in constructive logic, this is not so.

A sensible first question to ask might be: ‘do decidable propositions exist?’ Fortunately, the answer is affirmative.

- \top and \perp are decidable propositions;
- We would expect $m =_{\mathbb{N}} n$ to be a decidable proposition, where $=_{\mathbb{N}}$ denotes equality on the natural numbers;
- We would *not* expect $x =_{\mathbb{R}} y$ to be a decidable proposition, where $=_{\mathbb{R}}$ denotes equality on the real numbers, because real numbers are not finite objects.

Definition. A prop is *stable* if and only if $(\neg\neg A) \supset A \text{ true}$.

Again, in classical logic, every proposition is stable; in fact, the proposition $(\neg\neg A) \supset A \text{ true}$ is often taken as an axiom of treatments of classical propositional logic! A natural question to ask now is ‘do there exist unstable propositions?’ Consider the following lemma.

Lemma. $\neg\neg(A \vee \neg A) \text{ true}$

Proof. We must show $\neg(A \vee \neg A) \supset \perp \text{ true}$.

Suppose $A \text{ true}$. We then have

$$\frac{\frac{A \text{ true}}{A \vee \neg A \text{ true}} \vee I_1}{\perp} \neg(A \vee \neg A) \text{ true}$$

So in fact $\neg A \text{ true}$. But then once again

$$\frac{\frac{\neg A \text{ true}}{A \vee \neg A \text{ true}} \vee I_2}{\perp} \neg(A \vee \neg A) \text{ true}$$

Hence

$$\frac{\neg(A \vee \neg A) \text{ true} \vdash \perp}{\neg(A \vee \neg A) \supset \perp \text{ true}} \supset I$$

□

We can think of this lemma as saying that ‘the **law of the excluded middle** is not refutable’. Presuming that there exist undecidable propositions, we obtain the following corollary.

Corollary. In **intuitionistic propositional logic**, not every proposition is stable.

3 Disjunction property

A theory T has the **disjunction property (DP)** if $T \vdash A \vee B$ implies $T \vdash A$ or $T \vdash B$.

Theorem. In **IPL**, if $\emptyset \vdash A \vee B \text{ true}$ then $\emptyset \vdash A \text{ true}$ or $\emptyset \vdash B \text{ true}$.

Naïve attempt at proof. The idea is to perform induction on all possible derivations ∇ of $\emptyset \vdash A \vee B \text{ true}$, with the hope that somewhere along the line we’ll find a derivation of $A \text{ true}$ or of $B \text{ true}$. Our induction hypothesis is that inside ∇ is enough information to deduce either $\emptyset \vdash A \text{ true}$ or $\emptyset \vdash B \text{ true}$.

Since $\emptyset \vdash A \vee B \text{ true}$ cannot be obtained by assumption or from the rules, $\wedge I$, $\supset I$ or $\top I$, we need only consider $\vee I_1$, $\vee I_2$ and the elimination rules.

If $\emptyset \vdash A \vee B \text{ true}$ is obtained from $\vee I_1$ then

$$\frac{\frac{\nabla}{A \text{ true}}}{\emptyset \vdash A \vee B \text{ true}} \vee I_1$$

so there is a derivation ∇ of $A \text{ true}$ and we’re done. Likewise if $\emptyset \vdash A \vee B \text{ true}$ is obtained from $\vee I_2$ then there is a derivation of $B \text{ true}$.

If $\emptyset \vdash A \vee B \text{ true}$ is obtained from $\supset E$ then the deduction takes the form

$$\frac{\frac{\nabla_1}{\emptyset \vdash C \supset (A \vee B) \text{ true}} \quad \frac{\nabla_2}{\emptyset \vdash C \text{ true}}}{\emptyset \vdash A \vee B \text{ true}} \supset E$$

We (dubiously¹) assume that $\emptyset \vdash C \supset (A \vee B) \text{ true}$ must have been derived in some way from $C \text{ true} \vdash (A \vee B) \text{ true}$. Suppose that this happens and that ∇'_1 is a deduction

¹In fact, this ‘dubious’ assumption is true in constructive logic.

of $C \text{ true} \vdash (A \vee B) \text{ true}$. We can then ‘substitute’ ∇_2 for all the occurrences of the assumption $C \text{ true}$ appearing in ∇'_1 to obtain a smaller derivation ∇_3 of $\emptyset \vdash A \vee B \text{ true}$. Our induction hypothesis then gives us that inside ∇_3 is enough information to deduce $\emptyset \vdash A \text{ true}$ or $\emptyset \vdash B \text{ true}$.

A similar approach works (we hope) for $\wedge E$ and $\supset E$, thus giving the result. \square

4 Admissible properties

The sketch proof of the previous theorem relied on transitivity of \vdash ; namely, that the following rule is true:

$$\frac{\Gamma, A \text{ true} \vdash B \text{ true} \quad \Gamma \vdash A \text{ true}}{\Gamma \vdash B \text{ true}} \top$$

This leads us naturally into a discussion of the structural properties of \vdash .

Definition. A deduction rule is *admissible* (in **IPL**) if nothing changes when it is added to the existing rules of **IPL**.

To be clear about which logical system we use, we may write \vdash_{IPL} to denote deduction in **IPL** rather than in some new logical system.

The goal now is to prove that the structural rules for entailment (reflexivity, transitivity, weakening, contraction, exchange) are admissible.

Theorem. The structural properties of \vdash_{IPL} are admissible.

Proof. **R, C, X:** Reflexivity, contraction and exchange are all primitive notions, in that they follow instantly. For instance:

$$\frac{\frac{\Gamma \vdash A \text{ true}}{\Gamma \vdash A \wedge A \text{ true}} \wedge I}{\Gamma \vdash A \text{ true}} \wedge E_1$$

so if we were to introduce

$$\frac{\Gamma \vdash A \text{ true}}{\Gamma \vdash A \text{ true}} \text{R}$$

as a new rule, then nothing would change. (Likewise for contraction and exchange.)

W: For weakening we use the fact that the structural rules are *polymorphic* in Γ . We can thus prove that weakening is admissible by induction: if the following rules are admissible

$$\frac{\Gamma \vdash B_2 \text{ true}}{\Gamma, A \text{ true} \vdash B_1 \text{ true}} \quad \text{and} \quad \frac{\Gamma \vdash B_2 \text{ true}}{\Gamma, A \text{ true} \vdash B_1 \text{ true}}$$

then we obtain

$$\frac{\frac{\frac{\Gamma \vdash B_1 \wedge B_2 \text{ true}}{\Gamma \vdash B_1 \text{ true}} \wedge E_1 \quad \frac{\Gamma \vdash B_1 \wedge B_2 \text{ true}}{\Gamma \vdash B_2 \text{ true}} \wedge E_2}{\Gamma, A \text{ true} \vdash B_1 \text{ true}} \text{Ind} \quad \frac{\Gamma \vdash B_1 \wedge B_2 \text{ true}}{\Gamma, A \text{ true} \vdash B_2 \text{ true}} \text{Ind}}{\Gamma, A \text{ true} \vdash B_1 \wedge B_2 \text{ true}} \wedge I$$

Likewise for the other introduction rules.

⊤: The admissibility of transitivity is left as an exercise. \square

5 Proof Terms

We wish to study propositions along with their proof as mathematical objects. In the type theoretic framework, we can use the notation $M : A$ where A is a proposition and M is a proof of A . We will see that this corresponds to the category theoretic notion of a mapping $M : A \rightarrow B$. Another important notion is the identity of proofs, which will be denoted $M \equiv N : A$ where M, N are equivalent proofs of A . This will correspond in the category theoretic context to two maps from A to B being equal $M = N : A \rightarrow B$.

5.1 Proof Terms as Variables

We can combine the idea of keeping track of proofs with our previous notion of entailment. If A_1, \dots, A_n entails A , meaning that $A_1, \dots, A_n \vdash A$, there will be a proof M of A that uses the propositions A_1, \dots, A_n . Thus, we will write

$$x_1 : A_1, \dots, x_n : A_n \vdash M : A$$

where each $x_i : A_i$ is a proof term. We can think of the proof terms x_1, \dots, x_n as hypotheses for the proof, but what we really want is for them to behave as variables. M then uses the variables x_1, \dots, x_n to prove A , so M would encapsulate the grammar a proof that uses variables x_1, \dots, x_n .

Instead of proving a proposition A from nothing, most of the time A will rely on other propositions A_1, \dots, A_n .

5.2 Structural Properties of Entailment with Proof Terms

Now that we have proof terms, we can see how they act as variables by examining their interaction with the structural properties of entailment. We will also keep track of other assumptions/context Γ, Γ' to demonstrate that the structural properties will hold in the presence of assumptions.

Reflexivity / Variables Rule Reflexivity tells us that A should entail A , so now that we have a variable $x : A$ that proves A , the variable should be carried through. We can think of this as the variables rule.

$$\frac{}{\Gamma, x : A, \Gamma' \vdash x : A} \text{R/V}$$

Transitivity / Substitution Transitivity tells us that if A is true and B follows from A , then B is true. In terms of proofs, if we have a proof $N : A$ and a proof $N : B$ which uses a variable x that is supposed to prove A , then we can substitute the proof $M : A$ into $N : B$ to prove B . Since we are substituting M into x inside N , we denote this substitution $[M/x]N : B$.

$$\frac{\Gamma, x : A, \Gamma' \vdash N : B \quad \Gamma \vdash A}{\Gamma, \Gamma' \vdash [M/x]N : B} \text{T/S}$$

Weakening

$$\frac{\Gamma \vdash M : A}{\Gamma, \Gamma' \vdash M : A} \text{W}$$

Contraction If $N : B$ follows from A using two different proofs $x : A, y : A$ for A , can just pick one $z = x$ or $z = y$ as the proof of $z : A$ and use it in the instances of variables x, y in $N : B$

$$\frac{\Gamma, x : A, y : A, \Gamma' \vdash N : B}{\Gamma, z : A, \Gamma' \vdash [z, z/x, y]N : B} \text{C}$$

Exchange

$$\frac{\Gamma, x : A, y : B, \Gamma' \vdash N : C}{\Gamma, y : B, x : A, \Gamma' \vdash N : C} \text{X}$$

5.3 Negative Fragment of IPL with Proof Terms

We want to look at what happens to the Negative Fragment of IPL when we consider proof terms. Here are the important ones:

Truth Introduction Truth is trivially true, so we have

$$\frac{}{\Gamma \vdash \langle \rangle : \top} \text{T/I}$$

Conjunction Introduction We combine the proofs $M : A$ and $N : B$ into $\langle M, N \rangle : A \wedge B$

$$\frac{\Gamma \vdash M : A \quad \Gamma \vdash N : B}{\Gamma \vdash \langle M, N \rangle : A \wedge B} \wedge I$$

Conjunction Elimination We can recover from a proof $M : A \wedge B$ proofs of A and B

$$\frac{\Gamma \vdash M : A \wedge B}{\Gamma \vdash \text{fst}(M) : A} \wedge E_1 \qquad \frac{\Gamma \vdash M : A \wedge B}{\Gamma \vdash \text{snd}(M) : B} \wedge E_2$$

Implication Introduction If we have a proof $M : B$ that uses $x : A$ as a variable, then we can consider $\lambda x.M$ as a function that maps x a variable to a proof of B that uses x , which proves that $B \supset A$

$$\frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash \lambda x.M : A \supset B} \supset I$$

Implication Elimination By applying an actual proof $N : A$ to the function described above, we obtain a proof $M(N) : B$

$$\frac{\Gamma \vdash M : A \supset B \quad \Gamma \vdash N : A}{\Gamma \vdash M(N) : B} \supset E$$

6 Identity of Proofs

6.1 Definitional Equality

We want to think about when two proofs $M : A$ and $M' : A$ are the same. We will introduce an equivalence relation called *definitional equality* that respects the proof rules, denoted $M \equiv M' : A$. We want definitional equality \equiv to be the least congruence containing (closed under) the β rules. We will define what this means:

A *congruence* is an equivalence relation that respects our operators. Being an equivalence relation that it is reflexive ($M \equiv M : A$), symmetric ($M \equiv N : A$ implies that $N \equiv M : A$), and transitive ($M \equiv N : A$ and $N \equiv M' : A$ implies that $M \equiv M' : A$).

For the equivalence relation to respect our operators basically means that if $M \equiv M' : A$, then that their image under any operator should be equivalent. In

other words, we should be able to replace M with M' everywhere. For example

$$\frac{\Gamma \vdash M \equiv M' : A \wedge B}{\Gamma \vdash \text{fst}(M) \equiv \text{fst}(M') : A}$$

There can be many congruences that contains the β rules. Given two congruences \equiv and \equiv' , we say \equiv is finer than \equiv' if $M \equiv' N : A$ implies that $M \equiv N : A$. The least congruence that contains the proof rules is the finest congruence that contains the β rules. We will define the β rules in the next section.

We will give a more explicit definition to definitional equality later.

6.2 Gentzen's Inversion Principle

Gentzen's Inversion Principle captures the idea that “elim is post-inverse to intro,” which is the informal notion that the elimination rules should cancel the introduction rules, modulo definitional equality. The following are the β rules for the negative fragment of IPL:

Conjunction When we introduce a conjunction, we combine proofs $M : A$ and $N : B$ to produce a proof $\langle M, N \rangle : A \wedge B$. When we eliminate a conjunction, we retrieve $M : A$ or $N : B$. We do not want this process to alter our original M or N

$$\frac{\Gamma \vdash M : A \quad \Gamma \vdash N : B}{\Gamma \vdash \text{fst}(\langle M, N \rangle) \equiv M : A} \beta_{\wedge_1}$$

$$\frac{\Gamma \vdash M : A \quad \Gamma \vdash N : B}{\Gamma \vdash \text{snd}(\langle M, N \rangle) \equiv N : A} \beta_{\wedge_2}$$

Implication When we introduce an implication, we convert a proof $M : B$ which uses some variable $x : A$ to a function which uses a variable x to produce a proof of B . When we eliminate implication, we apply the proof of $A \supset B$ to $N : A$ to produce a proof of B .

$$\frac{\Gamma, x : A \vdash M : B \quad \Gamma \vdash N : A}{\Gamma \vdash (\lambda x. M)(N) \equiv [N/x]M : B} \beta_{\supset}$$

6.3 Gentzen's Unicity Principle

Gentzen's Unicity Principles on the other hand captures the idea that “intro is post-inverse to elim.” Another way to think about it is that there should be only one way modulo definitional equivalence to prove something, which is the way we have described. They are the η rules, which are the following

Truth

$$\frac{\Gamma \vdash M : \top}{\Gamma \vdash M \equiv \langle \rangle : \top} \eta^\top$$

Conjunction

$$\frac{\Gamma \vdash M : A \wedge B}{\Gamma \vdash M \equiv \langle \text{fst}(M), \text{snd}(N) \rangle : A \wedge B} \eta^\wedge$$

Implication

$$\frac{M : A \supset B}{\Gamma \vdash M \equiv \lambda x. Mx : A \supset B} \eta^\supset$$

7 Proposition as Types

There is a correspondence between propositions and types:

Propositions	Types
\top	1
$A \wedge B$	$A \times B$
$A \supset B$	function $A \rightarrow B$ or B^A
\perp	0
$A \vee B$	$A + B$

For now, note that meets like \top and $A \wedge B$ corresponds to products like 1 and $A \times B$, and joins like \perp and $A \vee B$ corresponds to coproducts like 0 and $A + B$. This correspondence should become more apparent as we go along. We will now introduce the objects on the right column.

8 Category Theoretic Approach

In a Heyting Algebra, we have a preorder $A \leq B$ when A implies B . However, we now wish to keep track of proofs, so if M is a proof from A to B , we want to think of it as a map $M : A \rightarrow B$.

Identity There should be an identity map

$$\text{id} : A \rightarrow A$$

Composition We should be able to compose maps

$$\frac{g : B \rightarrow C \quad f : A \rightarrow B}{f \circ g : A \rightarrow C}$$

Coherence Conditions The identity map and composition of maps should behave like functions

$$\begin{aligned} \text{id}_B \circ f &= f : A \rightarrow B \\ f \circ \text{id}_A &= f : A \rightarrow B \\ f \circ (g \circ h) &= (f \circ g) \circ h : A \rightarrow D \end{aligned}$$

Now we can think about objects in the category that corresponds to propositions given in the correspondence.

Terminal Object 1 is the terminal object, also called the final object, which corresponds to \top . For any object A there is a unique map $A \rightarrow 1$. This corresponds to \top being the the greatest object in a Heyting Algebra, meaning that for all A , $A \leq 1$).


Existence:

$$\langle \rangle : A \rightarrow 1$$

Uniqueness:

$$\frac{M : A \rightarrow 1}{M = \langle \rangle : A \rightarrow 1} \eta_{\top}$$

Product For any objects A and B there is an object $C = A \times B$ that is the *product* of A and B , which corresponds to the join $A \wedge B$. The product $A \times B$ has the following universal property:



where the diagram commutes.

First, the existence condition means that there are maps

$$\begin{aligned} \text{fst} &: A \times B \rightarrow A \\ \text{snd} &: A \times B \rightarrow B \end{aligned}$$

The universal property says that for every object D such that $M : D \rightarrow A$ and $N : D \rightarrow B$, there exists a unique map $\langle M, N \rangle : D \rightarrow A \times B$ such that

$$\frac{M : D \rightarrow A \quad N : D \rightarrow B}{\langle M, N \rangle : D \rightarrow A \times B}$$

and the diagram commutes meaning

$$\begin{aligned} \text{fst} \circ \langle M, N \rangle &= M : D \rightarrow A && (\beta \times_1) \\ \text{snd} \circ \langle M, N \rangle &= N : D \rightarrow B && (\beta \times_2) \end{aligned}$$

Furthermore, the map $\langle M, N \rangle : D \rightarrow A \times B$ is unique in the sense that

$$\frac{P : D \rightarrow A \times B \quad \text{fst} \circ P = M : D \rightarrow A \quad \text{snd} \circ P = N : D \rightarrow B}{P = \langle M, N \rangle : D \rightarrow A \times B} \eta \times$$

so in other words $\langle \text{fst} \circ P, \text{snd} \circ P \rangle = P$.

Another way to say the above is

$$\begin{aligned} \langle \text{fst}, \text{snd} \rangle &= \text{id} \\ \langle M, N \rangle \circ P &= \langle M \circ P, N \circ P \rangle \end{aligned}$$

Exponentials Given objects A and B , an exponential B^A (which corresponds to $A \supset B$) is an object with the following universal property:

$$\begin{array}{ccc} C & C \times A & \\ \downarrow \lambda(h) & \downarrow \lambda(h) \times \text{id}_A & \searrow h \\ B^A & B^A \times A & \xrightarrow{\text{ap}} B \end{array}$$

such that the diagram commutes.

This means that there exists a map $\text{ap} : B^A \times A \rightarrow B$ (application map) that corresponds to implication elimination.

The universal property is that for all object C that has a map $h : C \times A \rightarrow B$, there exists a unique map $\lambda(h) : C \rightarrow B^A$ such that

$$(\lambda(h) \times \text{id}_A) \circ \text{ap} = h : C \times A \rightarrow B$$

This means that the diagram commutes. Another way to express the induced map is $\lambda(h) \times \text{id}_A = \langle \lambda(h) \circ \text{fst}, \text{snd} \rangle$.

The map $\lambda(h) : C \rightarrow B^A$ is unique, meaning that

$$\frac{\text{ap} \circ (g \times \text{id}_A) = h : C \times A \rightarrow B}{g = \lambda(h) : C \rightarrow B^A} \eta$$

References